

# Galois correspondences

David Pierce

2004.12.16

This note gives a unified treatment to several mathematical observations.

For notation, let  $2$  be considered as the universe  $\{0, 1\}$  of an abelian group. For each  $e$  in  $2$ , let  $A_e$  be a set, and let  $*$  be a relation from  $A_0$  to  $A_1$ . This means that  $*$  is a subset of  $A_0 \times A_1$ ; if  $*$  contains  $(x_0, x_1)$ , then we write

$$x_0 * x_1.$$

If  $x_e \in A_e$ , let

$$[x_e] = \{x_{e+1} \in A_{e+1} : x_0 * x_1\} \in \mathcal{P}(A_{e+1}).$$

If  $X \in \mathcal{P}(A_e)$ , let

$$X' = \bigcap \{[x] : x \in X\} \in \mathcal{P}(A_{e+1}).$$

Let  $A_{e+1}^*$  be the image of  $\mathcal{P}(A_e)$  under the map  $X \mapsto X'$ .

**Theorem 1.** *Let  $e \in 2$ .*

- (\*) *If  $X, Y \in \mathcal{P}(A_e)$  and  $X \subseteq Y$ , then  $Y' \subseteq X'$ .*
- (†) *If  $X \in \mathcal{P}(A_e)$ , then  $X \subseteq X''$ .*
- (‡) *The map  $X \mapsto X'$  is a bijection from  $A_e^*$  to  $A_{e+1}^*$  with inverse  $X \mapsto X'$ .*

*Proof.* Exercise. (For the last point, see [3, ch. V, Lemma 2.6].) □

Regardless of how the maps  $X \mapsto X'$  are originally defined, if they meet the conditions established in the theorem, they constitute a **Galois correspondence** between  $A_0^*$  and  $A_1^*$ . (This definition is in [4, p. 35].) There are several examples, as you should verify:

## Field-theory

The usual Galois correspondence in field-theory is the case when  $A_0$  is a field  $L$  that is a finite Galois extension of a field  $K$ , and  $A_1$  is  $\text{Aut}(L/K)$ , and

$$x * \sigma \iff x^\sigma = x.$$

Then  $A_0^*$  comprises the subfields  $F$  of  $L$  that include  $K$ , and  $A_1^*$  comprises the subgroups  $H$  of  $\text{Aut}(L/K)$ , and  $F' = \text{Aut}(L/F)$ , and  $H' = \text{Fix}(H)$ .

## The Zariski topology

Suppose  $A_0$  is a ring  $R$  (commutative with 1), and  $A_1$  is  $\text{Spec } R$ , that is, the set of prime ideals of  $R$ . Let  $*$  be  $\in$ . Then

$$[x] \cup [y] = [xy]$$

if  $x, y \in R$ . Hence the sets  $[x]$  are the basic closed sets for a topology, the **Zariski topology** on  $\text{Spec } R$ . (See for example [1, pp. 54–55] or [2, § II.2].) The topology is compact, although possibly not Hausdorff. In this topology, if  $X \subseteq \text{Spec } R$ , then  $X''$  is the closure of  $X$ . If  $X \subseteq R$ , then  $X''$  is the radical (in the sense of [3, ch. VIII, Definition 2.5]) of the (possibly improper) ideal  $(X)$ . In general,  $A_0^*$  comprises the radical ideals of  $R$ , and  $A_1^*$  comprises the closed subsets of  $\text{Spec } R$ .

## The Stone space

Now suppose in particular that  $A_0$  is a Boolean ring or algebra  $B$ , and  $A_1$  is its **Stone space**  $S(B)$ , the set of ultrafilters of  $B$ . The ultrafilters are dual to the prime ideals, all of which are maximal. Let  $*$  be  $\in$  again. Then

$$[x] \cap [y] = [xy] = [x \wedge y]$$

when  $x, y \in B$ , and also

$$[x]^c = [x + 1] = [\neg x],$$

so that

$$[x] \cup [y] = [x + y + xy] = [x \vee y].$$

Hence the sets  $[x]$  are basic open and closed sets for a topology on  $S(B)$ . This topology is compact as before, but also Hausdorff. The elements of  $A_1^*$  are still just the closed subsets of  $S(B)$ ; the elements of  $A_0^*$  are just the filters of  $B$ . If  $X \subseteq B$ , then  $X''$  is the filter generated by  $X$ ; if  $X \subseteq S(B)$ , then  $X''$  is its closure.

## Model-theory

Suppose  $\mathcal{L}$  is a signature for first-order logic. Let  $A_0$  be the *class*  $\text{Mod}(\mathcal{L})$  of  $\mathcal{L}$ -structures, let  $A_1$  be  $\text{Sn}_{\mathcal{L}}$ , and let  $*$  be  $\models$ . Then  $A_1^*$  is the set of theories of  $\mathcal{L}$ , and  $A_0^*$  is the set of **elementary classes** of  $\mathcal{L}$ -structures. (See [4, § 3.4].)

## References

- [1] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [2] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.

- [3] Thomas W. Hungerford. *Algebra*, volume 73 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1980. Reprint of the 1974 original.
- [4] Philipp Rothmaler. *Introduction to model theory*, volume 15 of *Algebra, Logic and Applications*. Gordon and Breach Science Publishers, Amsterdam, 2000. Prepared by Frank Reitmaier, Translated and revised from the 1995 German original by the author.